

## REFERENCES:

1. Storm H. Skin conductance activity and the stress response from heel stick in premature infants. *Archives of Disease in Childhood* 2000;83(2):F143-F147.
2. Lidberg L, Wallin G. Sympathetic skin nerve discharges in relation to amplitude of skin resistance responses. *Psychophysiology* 1981;18(3):268-270.
3. Hagbarth KE, Hallin RG, Hongell A, Torebjørk HE, Wallin BG. General characteristics of sympathetic activity in human skin nerves. *Acta Physiologica Scandinavica*. 1972 (84):164-176.
4. Wallin BG, Sundløf G; Delius W. The effect of carotid sinus nerve stimulation on muscle and skin nerve sympathetic activity in man. *Pflügers Archiv* 1975(358):101-110.
5. Storm H, Myre K, Rostrup M, Stokland O, Ræder JC. A case report from a patient that wake up during surgery and was monitored by skin conductance and bispectral (BIS) index. *Minerva anesthesiologica* 2001(67)suppl 1 N 5:450.
6. Storm H, Fremming A, Ødegaard S, Martinsen Ø, Mørkrid L. A software analysis program for spontaneous and stimulated skin conductance activity. *Clinical Neurophysiology* 2000;111:1989-1998.
7. Storm H. The development of a software analysis program for analysing spontaneous skin conductance activity in preterm infants, *Clinical Neurophysiology* 2001(112):1562-1568.
8. Bini G, Hagbarth KE, Hynninen P, Wallin BG. Thermoregulatory and rythm-generating mechanisms govering the sudomotor and vasoconstrictor outflow in human cutaneous nerves.
9. Storm H, Myre K, Rostrup M, Stokland O, Ræder J. Skin conductance correlates with surgical stress and inversely with serum levels of general anaesthetics. *Minerva Anesthesiologica* 2001(67)suppl 1 N 5:153.
10. Hellerud BC, Wagner K, Henriksen T, Amo T, Storm H. Department of Pediatric Research and Department of Intensive Care, The National hospital, Department of Intensive Care, Ullevål University Hospital, Oslo, Institute of Pharmacology, Trondheim, Norway. Stress activation by suction from trackea in artificial ventilated children in intensive care units measured by skin conductance, heart rate, blood pressure, sedation score, the blood level of morphine, and the break down enzymes of morphine, in prep.
11. Storm H. A pilot study of how skin conductance changes may be used as a method to measure pain responses in patients with chronicle pain. *Journal of Psychosomatic Research* 2001: Abstract of the 24<sup>th</sup> ECPR.
12. Luis R, Storm H. Department of Paediatric Research and Department of Anaesthesiology, The National Hospital, Oslo. Measuring of stress and pain responses on voluntary adult persons after a burning injury by visual analogue scale, blood pressure, heart rate and skin conductance variables, in prep.
13. Brattebø G, Gjerde S, Muri AK, Aarland IC, Flaatten HK. Reducing ventilatory time in adult intensive care patients by optimising sedation practice. *Intensive care medicine* 2000;26:S384.
14. Eide PK, Storm H, Fremming A, Due-Tønnessen B, Lundar T. Department of Neuro surgery and Deparment of Pediatric Research, The National Hospital, Oslo. Spontaneous skin conductance as a sign of pain during increased intracranial pressure, in prep.

15. Storm H, Østberg BC. A pilot study of spontaneous and stimulated skin conductance changes in a patient suffering from anorexia nervosa. *Journal of Psychosomatic Research* 48 (2000): 230.

16. Storm H, Vandvik IH. A pilot study of how spontaneous skin conductance changes may be used as a method to measure the effect of hypnosis. *Journal of Psychosomatic Research* 48 (2000):276.

17. Jørum E, Storm H et al. Injury in the peripheral nervous system and skin conductance changes, in preparation.